Conference Abstract

OPEN

Application of geomicrobial techniques to constrain mechanisms of arsenic mobilisation in anoxic aquifers

Teto I Seitshiro[‡], Naji M Bassil[‡], Oliver Moore[‡], Jonathan R Lloyd[‡]

‡ The University of Manchester, Manchester, United Kingdom

Corresponding author: Teto I Seitshiro (roman.seitshiro@manchester.ac.uk)

Received: 09 Jun 2023 | Published: 13 Oct 2023

Citation: Seitshiro TI, Bassil NM, Moore O, Lloyd JR (2023) Application of geomicrobial techniques to constrain mechanisms of arsenic mobilisation in anoxic aquifers. ARPHA Conference Abstracts 6: e107756. https://doi.org/10.3897/aca.6.e107756

Abstract

Geogenic arsenic contamination of groundwater in South and South-East Asia poses a significant human health threat, causing a range of health conditions including but not limited to cardio-vascular disease, cancer and skin lesions (Argos et al. 2010, Pienkowska et al. 2021). Arsenic contamination also hosts a range of dire socioeconomic implications for the affected areas. A variety of mechanisms for arsenic release in anoxic aquifers have been proposed, however, the most widely accepted mechanism is the microbial reduction of As-bearing Fe(III) (oxyhydr)oxide minerals coupled with the oxidation of organic carbon (Glodowska et al. 2020, Gnanaprakasam et al. 2017). Recent research has implicated methane as a possible carbon source in the reduction of Fe(III) (oxyhydr)oxide minerals and the subsequent release of arsenic into the groundwater (Gnanaprakasam et al. 2017, Pienkowska et al. 2021). The research suggests that methanotrophs have the ability to drive anaerobic oxidation of methane, AOM, coupled to Fe(III) (oxyhydr)oxide reduction. In this study, we aim to provide unequivocal evidence for the occurrence of AOM coupled to Fe(III) (oxyhydr)oxide as well to further explore the exact mechanism(s) involved which is yet to be characterised. Here we present an overview of our work so far.

© Seitshiro T et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords

Fe(III) (oxyhydr)oxides, Arsenic, Methane

Presenting author

Teto Seitshiro

Funding program

Botswana Government Top Achiever's Sholarship

Hosting institution

The University of Manchester

Conflicts of interest

The authors have declared that no competing interests exist.

References

- Argos M, Kalra T, Rathouz PJ, Chen Y, Pierce B, Parvez F, Islam T, Ahmed A, Rakibuz-Zaman M, Hasan R, Sarwar G, Slavkovich V, van Geen A, Graziano J, Ahsan H (2010) Arsenic exposure from drinking water, and all-cause and chronic-disease mortalities in Bangladesh (HEALS): a prospective cohort study. The Lancet 376 (9737): 252-258. https://doi.org/10.1016/s0140-6736(10)60481-3
- Glodowska M, Stopelli E, Schneider M, Rathi B, Straub D, Lightfoot A, Kipfer R, Berg M, Jetten MSM, Kleindienst S, Kappler A (2020) Arsenic mobilization by anaerobic irondependent methane oxidation. PANGAEA <u>https://doi.org/10.1594/pangaea.924771</u>
- Gnanaprakasam E, Lloyd J, Boothman C, Ahmed KM, Choudhury I, Bostick B, van Geen A, Mailloux B (2017) Microbial Community Structure and Arsenic Biogeochemistry in Two Arsenic-Impacted Aquifers in Bangladesh. mBio 8 (6). <u>https://doi.org/10.1128/</u> mbio.01326-17
- Pienkowska A, Glodowska M, Mansor M, Buchner D, Straub D, Kleindienst S, Kappler A (2021) Isotopic Labeling Reveals Microbial Methane Oxidation Coupled to Fe(III) Mineral Reduction in Sediments from an As-Contaminated Aquifer. Environmental Science & Technology Letters 8 (9): 832-837. <u>https://doi.org/10.1021/acs.estlett.</u> <u>1c00553</u>